arXiv:0805.2237v1 [hep-ph] 15 May 2008

Complete Gluonic Phase in Two-Flavor Color Superconductivity [*]

Xiao-Ming Wang[†] and Bang-Rong Zhou[‡]
College of Physical Sciences, Graduate University of Chinese Academy of Sciences
Beijing 100049, China

Abstract

We study a typical complete gluonic phase (LGP) in two-flavor color superconductivity (2SC) by calculating the essential cubic and quartic interfering term between the gluonic condensates $\langle A_z^{(8)}\rangle$ and $\langle A_z^{(6)}\rangle$ with a gauged NJL model. It is proven that the coefficient of the cubic interfering term and the vacuum contributions of the cubic and quartic interfering term are all equal to zeroes. The coefficients of the quartic interfering term and the $\langle A_z^{(6)}\rangle$'s quartic self-interaction term at stationary points of LOFF phase are calculated. Comparisons among the effective potentials of LGP, g2SC and LOFF phase indicate that LGP could be the genuine ground state of 2SC for some reasonable parameters.

PACS numbers: 12.38.-t, 11.30.Qc, 26.60.+c, 11.15.Ex
Key Words: effective potential, gluonic phase, LOFF phase

Since 1998, color superconductivity (CSC) in dense quark matter has been a popular research topic. It has not only important theoretical significance for understanding the phase structure of Quantum Chromodynamics but also direct practical applications to exploring the inner cores of compact stars. In recent years, discovery of the chromomagnetic instability(CMI) [1] triggers another research tide, and many works have been done to cure the CMI problem.[5, 6, 7]. In all the present CMI-resolving schemes, the gluonic phase (GP) is naturally of much interest [2, 3]. However, the gluonic phase discussed so far [2, 5, 6, 7] can not be called complete gluonic phase, because the gluonic condensate $\langle A_z^{(8)}\rangle$ was not included in the previous calculations. Since the gluon field $A_z^{(8)}$ itself has also the problem of CMI, the genuine ground state must contain the parameter $\langle A_z^{(8)}\rangle$. The authors of the paper [4] put forward some genuine complete gluonic phase schemes including $\langle A_z^{(8)}\rangle$ (so called phase A-phase F in Ref.[4]). But up to now, no detailed research has been done on their stability.

It should be indicated that introduction of$\langle A_z^{(8)}\rangle$ in 2SC is equivalent to including the LOFF phase extensively studied so far, with the correspondence that the parameter of single plane wave LOFF state $q \rightarrow \frac{g}{2\sqrt{3}}\langle A_z^{(8)}\rangle$. Since the effective potential of LOFF phase has been calculated precisely and some simple analytic expressions for most of the other GP's parameters have been obtained [2], so calculating the interfering terms between $\langle A_z^{(8)}\rangle$ and the other parameters of the GP becomes the key to derive the effective potential of a complete gluonic phase.

As stated above, introduction of the $\langle A_z^{8}\rangle$ is equivalent to inclusion of LOFF phase. However, in weak coupling region, LOFF phase might well be the ground state of 2SC and if that is true, then introduction of the other gluonic condensates will be unnecessary, because in this case LOFF phase itself can cure the CMI problem [8]. Otherwise, it will also be necessary to include the gluonic condensates other than $\langle A_z^{(8)}\rangle$. On the other hand, in the intermediate and strong coupling region, it has been proven that the ground state of 2SC can not be LOFF phase [9], hence the left alternative could only be the complete gluonic phase. For exploring such a phase, besides $\langle A_z^{8}\rangle$, we must at least include $\langle A_z^{(6)}\rangle$ which is the remaining non-zero component of a complex $SU_c(2)$-doublet composed of the gluonic condensates from $\vec{A}^{(4)} - \vec{A}^{(7)}$ in the unitary gauge and after an $O(3)$ rotation (similarly, $\langle A_z^{8}\rangle$ is the remainder of $\langle \vec{A}^{8}\rangle$ after an $O(3)$ rotation), here the $SU_c(2)$ is the remaining symmetry after the $SU_c(3)$ is broken by the two-flavor diquark condensates. The other condensates from the components $\langle \vec{A}^{(i)}\rangle (i = 1, 2, 3)$, though possible in principle, will not be considered in present model.

In this way, we will explore a typical edition of the complete gluonic phase (abbreviated as LGP, implying that the LOFF phase with $\langle A_z^{(8)}\rangle$ and the GP with $\langle A_z^{(6)}\rangle$ coexist). Let T_{Bq} and λ_{Bq} denote respectively the coefficient of the cubic and quartic interfering term between $\langle A_z^{(8)}\rangle$ and $B = \langle A_z^{(6)}\rangle$,

---

[*] The project supported by the National Natural Science Foundation of China under Grant No.10475113.

[†] E-mail:xiaomingw@mails.gucas.ac.cn
[‡] E-mail:zhoubr@gucas.ac.cn

and λ_B denote the coefficient of the quartic self-interaction term of B. We will only calculate the values of these coefficients and the LGP effective potential at the LOFF phase's ground state so as to examine the essential character of LGP. This is because the interfering terms between $\langle A_z^{(8)}\rangle$ and $B = \langle A_z^{(6)}\rangle$ haven't simple analytic expressions, so it is difficult to find out the solutions of the neutral condition and the gap equations and to fix the ground state's parameters of LGP. Hence, we have to adopt a conventional practice instead, i.e. takeing the ground state's parameters of other approximate phase as the ones of present model. It is indicated that the ground state's parameters of LOFF phase are close to the ones of LGP. In fact, the main part in the effective potential of LGP comes from 2SC+LOFF phase, thus the ground state's parameters of LGP will be mainly determined by the 2SC+LOFF sector in LGP. This point has been verified indirectly by the results in Refs. [5, 6]. For example, the parameters of the gluonic cylindrical phase II's ground state shown in **Fig.1** and **Fig.2** of Ref. [6] is surprisingly similar to those of the LOFF phase'ground state obtained by us and the small difference between the both may be attributed to different approximations to the effective potentials. Thus it is certainly rational to calculate the effective potential of LGP at the stationary points of LOFF phase. In addition, we can see from Refs.[5, 6] that the gluonic phase only gives no more than ten percent contribution to the total effective potential of 2SC +GP, hence in the Ginzburg-Landzu approach, the effective potential of LGP will be expanded in the gluonic condensates only up to the fourth order.

We will use a gauged NJL model describing two-flavor color superconductivity at zero temperature. When there exist diquark condensates, we introduce the Nambu-Gor'kov bispinor field

$$\Psi = \frac{1}{\sqrt{2}}\begin{pmatrix} q \\ q^C \end{pmatrix} \tag{1}$$

with the quark field $q \equiv q_{i\alpha}$ which is a four-component Dirac spinor with the flavor($i = u, d$) and the color ($\alpha = r, g, b$) indices and the charge conjugated field defined by $q^C = C\overline{q}^T$. Thus the Lagrangian density can be written as

$$\mathcal{L} = \overline{\Psi}\mathcal{S}^{-1}\Psi, \tag{2}$$

where the inverse propagator of the field Ψ

$$\mathcal{S}^{-1} = \begin{pmatrix} [G_0^+]^{-1} & \Delta^{-1} \\ \Delta^+ & [G_0^-]^{-1} \end{pmatrix} \tag{3}$$

with

$$\begin{aligned} [G_0^+]^{-1}(K) &= (k_0 + \overline{\mu} - \delta\mu\tau_3 - \mu_8\mathbf{1}_b)\gamma^0 \\ &\quad -\vec{\gamma}\cdot\vec{k}, & (4)\\ [G_0^-]^{-1}(K) &= (k_0 - \overline{\mu} + \delta\mu\tau_3 + \mu_8\mathbf{1}_b)\gamma^0 \\ &\quad -\vec{\gamma}\cdot\vec{k}, & (5)\\ \Delta^- &= -\imath\varepsilon\varepsilon^b\gamma_5\Delta, \\ \Delta^+ &= \gamma^0(\Delta^-)^\dagger\gamma^0 = -\imath\varepsilon\varepsilon^b\gamma_5\Delta & (6)\end{aligned}$$

The denotations ε, ε^b, $\overline{\mu}$, $\delta\mu$, μ_8, etc. in formulas (4)-(6) are the same as the ones used in Ref. [4].

The propagator of Ψ is given by

$$\begin{pmatrix} G^+ & \Xi^- \\ \Xi^+ & G^- \end{pmatrix} \tag{7}$$

with $G^\pm = ([G_0^\pm]^{-1} - \Delta^\mp G_0^\mp \Delta^\pm)^{-1}$, $\Xi^\pm = -G_0^\mp\Delta^\pm G^\pm$.

In order to cure CMI problem, we must extend S^{-1} to including the VEVs of relevant gluon fields. The effective potential of LGP in the Ginzburg-Landau approach can be generally expressed by [4]

$$\begin{aligned} V_{LGP} &= \frac{\Delta^2}{4G_D} - \frac{1}{2}\int\frac{d^4k}{i(2\pi)^4}Tr\mathcal{S}_g^{-1} \\ &\quad +\frac{1}{2}\int\frac{d^4k}{i(2\pi)^4}Tr\mathcal{S}_g^{-1}|_{\mu=\mu_e=\Delta=0} \\ &= V_\Delta + \Sigma_{n=1}^{\infty}\frac{(-1)^n}{2n}\int\frac{d^4k}{i(2\pi)^4}Tr(\mathcal{S}\mathcal{M}_g)^n \\ &\quad +\frac{1}{2}\int\frac{d^4k}{i(2\pi)^4}Tr\mathcal{S}_g^{-1}|_{\mu=\mu_e=\Delta=0}, \end{aligned} \tag{8}$$

where V_Δ represents the effective potential of 2SC and $\mathcal{S}_g^{-1} = S^{-1} + \mathcal{M}_g$ with $\mathcal{M}_g$ denoting the inverse propagator of relevant gluon fields. In present model, as has been stated above, only the two most important VEVs of gluon fields $\langle A_z^{(8)}\rangle$ and $B = \langle A_z^{(6)}\rangle$ will be included in $\mathcal{M}_g$. We will also use a simplifying treatment, i.e. setting $\mu_8 = 0$, as was done in other papers. On these grounds, through a lengthy calculation of relevant one-loop diagrams from the fermion determinant, we have got the expressions of T_{Bq} and λ_{Bq} in the form of products of fermion propagators. The the last term in (8) contains the vacuum contributions of the cubic and quartic interfering term between $\langle A_z^{(8)}\rangle$ and $B = \langle A_z^{(6)}\rangle$ which, however, are eventually proven to be equal to zeroes after a hard and careful calculation. As a result, the derived effective potential becomes

$$\begin{aligned} V_{LGP} &= V_\Delta + V_{LOFF} + M_B^2B^2/2 + \lambda_B B^4/4 \\ &\quad +T_{Bq}B^2q + \lambda_{Bq}B^2q^2/2, \end{aligned} \tag{9}$$

where V_{LOFF} is the effective potential of LOFF phase, M_B^2 is the squared Meissner mass of the gluonic field $A_z^{(6)}$. Furthermore, it has been proven that the coefficient of the cubic interfering term is equal to zero, i.e. $T_{Bq} = 0$. Hence the extreme value condition about parameter B derived from (9) becomes

$$\partial V_{LGP}/\partial B = M_B^2 B + \lambda_B B^3 + \lambda_{Bq} B q^2 = 0. \quad (10)$$

The solution of (10) is $B = 0$ and $B = B_1 = \sqrt{(-M_B^2 - \lambda_{Bq} q^2)/\lambda_B}$. It is seen from (9) that $B = 0$ just corresponds to the 2SC+ LOFF phase. By analyzing the second-order derivative of V_{LGP} over B, we find that if $\lambda_{Bq} < -M_B^2/q^2$, then $B = B_1$ and $B = 0$ will respectively be a minimal and a maximal point of V_{LGP} about B and this means that LGP could have lower energy than 2SC + LOFF phase in this case. Now considering only the situation of $B \neq 0$ and substituting B_1 into the effective potential (9), we may change V_{LGP} into $\bar{V}_{LGP}$, where

$$\begin{aligned} \bar{V}_{LGP} &= V_\Delta + V_{LOFF} - \lambda_B B^4/4 \\ &= V_\Delta + V_{LOFF} \\ &\quad -(-M_B^2 - \lambda_{Bq} q^2)^2/4\lambda_B. \quad (11) \end{aligned}$$

$\bar{V}_{LGP}$ is the effective potential of LGP at its non-zero stationary point about the parameter B and will become our starting point to make a comparison among the effective potentials of LGP, g2SC and LOFF phase. In fact, from (11), we can see that if $\lambda_B > 0$, then$\bar{V}_{LGP}$ could be lower than $V_\Delta + V_{LOFF}$.

The above qualitative analysis indicates that for making a comparison among the effective potentials, the key point is to calculate λ_{Bq} and λ_B. We find that in doing so, the most difficult task is to remove some superficial singularities. This is because after the energy summing, in the denominators of the integrands there will appear the factors similar to $a = p^0 - \sqrt{(|\vec{k}| + \bar{\mu})^2 + \Delta^2} + \sqrt{(|\vec{k}| + |\vec{p}| + \bar{\mu})^2 + \Delta^2}$ which could become zero when the gluons' four-momentum $(p^0, \vec{p}) \rightarrow 0$ and this will lead to singularity in the integral over the quark momentum $|\vec{k}|$. To remove such singularities, we invent a symbol calculating method. First denote these pole factors respectively by the symbols a,b,c,d,e,f and express an integrand generally in the form $f(a, b, c, d, e, f)/abcdef$, then by means of some found interrelations between these symbols, through lengthy calculations (since the quark propagators have large number -four- and very complicated forms), we have ultimately proven that all the pole factors in the denominators can be canceled out one by one and the superficial singularities are removed.



Fig.1: Varying of λ_{Bq} and λ_B as LOFF phase's stationary point when $\delta\mu > \Delta$ ($\mu = 0.4$ GeV, $\Lambda = 0.6533$ GeV).



Fig.2: Varying of λ_{Bq} and λ_B as LOFF phase's stationary point when $\delta\mu < \Delta$ ($\mu = 0.4$ GeV, $\Lambda = 0.6533$ GeV).

In our work, another key calculation is to fix the parameters of the ground state of LOFF phase. Although this calculation has been worked out [9], the detail of calculation wasn't given. In fact we have conducted this calculation independently and obtained the exactly same results as those shown in Ref. [9]. This lays a solid foundation for our following move. The last difficult task is to do the momentum integrations involving tedious numerical computations. The final results are shown in Fig.1, Fig.2 and Fig.3

All the x-axes of the three figures are Δ_0 which is the gap of the 2SC with $\delta\mu = 0$ and one-one corresponding to the coupling strength G_D of diquark condensates. For a fixed Δ_0 (G_D) we may find out a stationary point of LOFF phase (when μ is given), and in the meantime, a respective definite value of λ_{Bq}, λ_B and the effective potentials, while different Δ_0 may correspond to different parameters $\delta\mu$ and Δ. Based on these data, we will be able to compare the least energy states of LGP, g2SC and LOFF phase.

From Fig.1, it may be seen that at the stationary point of LOFF phase when $\delta\mu > \Delta$, we have



Fig.3: Effective potential of LGP, g2SC and LOFF phase at LOFF's stationary points ($\mu = 0.4$ Gev, $\Lambda = 0.6533$ Gev).

$\lambda_{Bq} > 0$ (dot line) and $\lambda_B < 0$ (dot-dash line). The value of λ_B is very small in the region $\Delta_0 < 0.10269$ GeV (e.g. $\Delta_0 = 0.081225$ GeV, $\lambda_B \approx 0.018$ and $\Delta_0 = 0.10269$ GeV, $\lambda_B \approx 0.24$). Near the critical point $\delta\mu = \Delta$ ($\Delta_0 = 0.109683$ GeV), the absolute values of λ_{Bq} and λ_B become very large, this attributes to some terms containing the δ function $\delta\left(\delta\mu - \sqrt{(|\vec{k}| - \bar{\mu})^2 + \Delta^2}\right)$ in λ_{Bq} and λ_B which become very large near the critical point after the momentum integral. In view of (11), the above result shows that when $\delta\mu > \Delta$, the effective potential of LGP is higher than that of LOFF phase.

From Fig.2, we can see that at the stationary point of LOFF phase when $\delta\mu \leq \Delta$ (0.109683 GeV $\leq \Delta_0 \lesssim 0.137$ GeV), the result is that $\lambda_{Bq} < 0$ (0.109683 GeV$< \Delta_0 \lesssim 0.134$ GeV) (dot line) and $\lambda_{Bq} > 0$(0.134 GeV $\lesssim \Delta_0 \lesssim 0.137$ GeV) and $\lambda_B > 0$ (dash line).

In Fig.3, we have drawn the effective potentials of LGP, g2SC and LOFF phase. Here we only give the effective potential of LGP when $\delta\mu < \Delta$ and omit the one when $\delta\mu > \Delta$, since in the latter case LGP has higher effective potential than LOFF phase at the stationary point of LOFF phase. From Fig.3, we can see that the effective potential of LOFF phase (dot line) is lower than that of the g2SC phase(dash line) in the whole LOFF phase window 0.065 GeV $< \Delta_0 < 0.137$ GeV. However, LGP (dot-dash line) has lower effective potential than LOFF phase at stationary point of LOFF phase when $\delta\mu < \Delta$ (0.10968 GeV $< \Delta_0 < 0.137$ GeV). Hence, we may conclude that in this region, LGP could be the ground state of the system.

In Fig.3, the abnormal part in the effective potential of LGP, namely, from point A to point B, is caused by the sudden vanishing of the terms containing δ function $\delta\left(\delta\mu - \sqrt{(|\vec{k}| - \overline{\mu})^2 + \Delta^2}\right)$. With regard to the situation at the LOFF phase's stationary point when $\delta\mu > \Delta$ i.e. in weak coupling region, the genuine ground state could be either LOFF state, or another phase when other parameters are considered.

The above study of LGP shows that the complete gluonic phase indeed possesses the character of the genuine ground state for some reasonable parameters. Although present work is preliminary and remains to be improved further, it is believed that the derived qualitative conclusion will not be changed essentially.

References

[1] M.Huang and I.A.Shovkovy, 2004 *Phys. Rev.*D **70** 051501(R); 2004 *Phys. Rev.* D **70** 094030.

[2] E.V.Gorbar, M.Hashimoto and V.A.Miransky, 2006 *Phys. Lett.* B **632**, 305-312.

[3] V.P.Gusynin, V.A.Miransky and I.A. Shovkovy, 2004 *Phys. Lett.* B **581**, 82; 2004 *Mod. Phys. Lett.* A **19**, 1341.

[4] E.V.Gorbar, M.Hashimoto and V.A.Miransky, 2007 *Phys. Rev. D **75** 085012.*

[5] O.Kiriyama, D.H.Rischke and I.A.Shovkovy, 2006 *Phys. Lett.* B **643**, 331.

[6] M.Hashimoto and V.A.Miransky, 2007 *Prog. Theor. Phys. **118** 303.*

[7] K.Fukushima, 2006 *Phys. Rev.* D **73** 094016; O.Kiriyama, 2006 *Phys. Rev.* D **74** 074019; 114011.

[8] I.Ginnakis, De-Fou Hou and Hai-Chang Ren, 2005 *Phys. Lett.* B **631** 16.

[9] E.V.Gorbar, M.Hashimoto and V.A.Miransky, 2006 *Phys. Rev. Lett* **96**, 022005.